

QCD corrections to neutralino annihilation

Heather E. Logan

Ottawa-Carleton Institute for Physics, Carleton University, Ottawa K1S 5B6 Canada

Abstract. We report on two recent calculations of QCD corrections to neutralino annihilation cross sections: (1) the next-to-leading order corrections to $\chi\chi \rightarrow gg$, and (2) the contribution to the cross section for $\chi\chi \rightarrow q\bar{q}g$ arising from interference between the tree-level and loop-induced processes.

The presence of non-baryonic dark matter in the universe is compelling evidence for physics beyond the Standard Model. Supersymmetry (SUSY) provides an especially attractive explanation with the lightest supersymmetric particle, usually a neutralino χ , as the dark matter candidate. In this talk we review two recent calculations [1, 2] of QCD corrections to neutralino annihilation via internal squark exchange.

The tree-level neutralino annihilation process $\chi\chi \rightarrow q\bar{q}$ via exchange of a squark of mass $M_{\tilde{q}}$ can be reduced to an effective vertex described by a dimension-six operator suppressed by $M_{\tilde{q}}^2$,

$$\mathcal{L} = (c/M_{\tilde{q}}^2) \mathcal{O}_6, \quad \mathcal{O}_6 = (\bar{\chi}\gamma_\mu\gamma_5\chi)(\bar{q}\gamma^\mu\gamma_5q). \quad (1)$$

If the annihilation occurs at rest, the Majorana neutralino spinors reduce to a spin-singlet combination, $\bar{\chi}\gamma_\mu\gamma_5\chi \rightarrow K_\mu\bar{\chi}(\gamma_5/\sqrt{2}m_\chi)\chi$, where K is the center-of-mass momentum of the two-neutralino system. The operator $\mathcal{O}_6$ can then be written as the divergence of the axial-vector current,

$$\mathcal{O}_6 \rightarrow \left[\bar{\chi} (i\gamma_5/\sqrt{2}m_\chi)\chi \right] \left[\partial_\mu (\bar{q}\gamma^\mu\gamma_5q) \right]. \quad (2)$$

In the massless quark limit, $m_q = 0$, the axial vector current is conserved at tree level, $\partial_\mu (\bar{q}\gamma^\mu\gamma_5q) = 0$, and all tree-level dimension-six amplitudes vanish in this limit. This is the well known partially-conserved axial current (PCAC) condition.

The quark mass suppression can be lifted in two ways:

1. By going beyond leading order in α_s to include the correction to the dimension-six operator that involves the anomalous triangle diagram. This is $\chi\chi \rightarrow gg$. This process first appears at order α_s^2 , and thus suffers from a large QCD scale uncertainty. We compute the next-to-leading order (NLO) QCD corrections to this process [1].
2. By going to dimension-eight by including hard gluon radiation. This is $\chi\chi \rightarrow q\bar{q}g$. The tree-level diagram interferes with $\chi\chi \rightarrow gg^*$ with the off-shell gluon splitting into a $q\bar{q}$ pair, yielding a contribution to the cross section of order α_s^2 , the same order as the leading $\chi\chi \rightarrow gg$ cross section. We compute this interference term [2].

$\chi\chi \rightarrow gg$ at NLO: The leading-order (LO) amplitude for $\chi\chi \rightarrow gg$ in the massless quark limit is given by

$$\mathcal{M}_{\text{loop}} = \frac{\alpha_s}{2\sqrt{2}\pi} \sum_{q'} \left[\frac{|g_r|^2}{M_{\tilde{q}_R}^2} + \frac{|g_\ell|^2}{M_{\tilde{q}_L}^2} \right] i\varepsilon^{\mu\nu\alpha\beta} \varepsilon_\mu^{*a}(k_1) \varepsilon_\nu^{*a}(k_2) k_{1\alpha} k_{2\beta}, \quad (3)$$

where the sum runs over the five light (massless) quark flavors q' and we neglect terms subleading in m_χ^2/M_q^2 . The $\chi q' \tilde{q}_{R,L}'$ couplings $g_{r,\ell}$ for right and left quark helicities, respectively, are

$$g_r = -\sqrt{2}N_{11}g'Q, \quad g_\ell = -\sqrt{2}N_{11}g'(T_3 - Q) + \sqrt{2}N_{12}gT_3. \quad (4)$$

Here T_3 is the quark isospin, Q is the quark electric charge, g and g' are the weak couplings under $SU(2)_L$ and $U(1)_Y$, respectively, and N_{11} and N_{12} are the bino and wino components of the neutralino as defined in Ref. [3]. The matrix element in Eq. (3) results in an annihilation cross section for $\chi\chi \rightarrow gg$ of

$$v_{\text{rel}} \sigma_{\text{dim6}} = \frac{\alpha_s^2 m_\chi^2}{32\pi^3} \left\{ \sum_{q'} \left[\frac{|g_r|^2}{M_{\tilde{q}_R}^2} + \frac{|g_\ell|^2}{M_{\tilde{q}_L}^2} \right] \right\}^2. \quad (5)$$

For $6 \text{ GeV} < m_\chi < 110 \text{ GeV}$, this approximate formula is within 10% of the result including the full quark mass dependence [1].

In order to compute the NLO QCD corrections to $\chi\chi \rightarrow gg$, we exploit the anomaly equation to relate this process to pseudoscalar decay to gluons. The conservation of the axial vector current is violated by the quark mass and by the anomalous triangle diagram, in the form [4]

$$\partial_\mu (\bar{q} \gamma^\mu \gamma_5 q) = 2m_q \bar{q} i \gamma_5 q + \frac{\alpha_s}{4\pi} G_{\mu\nu}^{(a)} \tilde{G}^{(a)\mu\nu}, \quad (6)$$

with $\frac{1}{2} \tilde{G}_{\mu\nu} = \varepsilon_{\mu\nu\alpha\beta} G^{\alpha\beta}$ denoting the dual color field strength tensor.

The amplitude for $\chi\chi \rightarrow gg$ is obtained by setting $m_q = 0$ and neglecting the first term on the right hand side of Eq. (6). The gluonic decay amplitude of a fundamental pseudoscalar, $A^0 \rightarrow gg$, is also related to the anomaly equation. This decay proceeds through a quark loop; if the Yukawa coupling of this quark to A^0 is proportional to the quark mass, the decay is dominated by the contribution of the heaviest quark in the loop. In the heavy quark limit, $m_Q \gg m_A$, we can neglect the divergence term on the left-hand side of Eq. (6). Thus we see that two seemingly different processes, $A^0 \rightarrow gg$ through a heavy quark loop and $\chi\chi \rightarrow gg$ through a light quark loop, are related by the anomaly equation to the *same* gluonic operator. The Adler-Bardeen theorem [5] guarantees that Eq. (6) is valid to all orders in α_s . We take advantage of this to obtain the NLO QCD corrections to $\chi\chi \rightarrow gg$ from the known results for $A^0 \rightarrow gg$ at NLO.

In the heavy top quark limit, for which our anomaly relation is valid, the NLO QCD corrections to the $A^0 \rightarrow gg$ partial width are given by a multiplicative factor times the LO decay rate [6]. Invoking the Adler-Bardeen theorem, the NLO cross section for $\chi\chi \rightarrow gg$

in the zero-velocity limit becomes [1]

$$v_{\text{rel}} \sigma_{\text{NLO}}(\chi\chi \rightarrow gg) = v_{\text{rel}} \sigma_{\text{dim6}} \left[1 + \frac{\alpha_s}{\pi} \left(\frac{97}{4} - \frac{7}{6} N_f + \frac{33 - 2N_f}{6} \log \frac{\mu^2}{4m_\chi^2} \right) \right], \quad (7)$$

where μ is the renormalization scale. The strong coupling is evaluated based on N_f -flavor running at the scale μ where it appears both explicitly in the correction term and within the leading-order cross section $v_{\text{rel}} \sigma_{\text{dim6}}$. The integer N_f counts the number of quark flavors in the gluon splitting, with $N_f = 5$ for $m_b \ll m_\chi \ll m_t$. Taking $\mu = 2m_\chi$, $N_f = 5$, and $m_\chi = 100$ GeV, we obtain $v_{\text{rel}} \sigma_{\text{NLO}}(\chi\chi \rightarrow gg) \simeq v_{\text{rel}} \sigma_{\text{dim6}} [1 + 0.62]$. The NLO corrections reduce the residual renormalization scale dependence of the $\chi\chi \rightarrow gg$ annihilation cross section from $\pm 16\%$ to $\pm 9\%$ when we vary μ by a factor of two in either direction [1].

$\chi\chi \rightarrow q\bar{q}g$ interference term: To leading order in m_χ^2/M_q^2 , it is straightforward to extend the loop amplitude for $\chi\chi \rightarrow gg$ to include one off-shell gluon splitting into a quark-antiquark pair:

$$\mathcal{M}_{\text{split}} = -\frac{g_s}{\sqrt{2}} \sum_{q'} \left[\frac{|g_r|^2}{M_{\tilde{q}_R}^2} + \frac{|g_\ell|^2}{M_{\tilde{q}_L}^2} \right] i\varepsilon^{\mu\nu\alpha\beta} \varepsilon_\mu^{*c} q_{3\alpha} (q_1 + q_2)_\beta \frac{\alpha_s \bar{u}(q_1) T^c \gamma_\nu v(q_2)}{2\pi (q_1 + q_2)^2}, \quad (8)$$

where q_1 , q_2 , and q_3 are the momenta of the final-state quark, antiquark, and gluon, respectively, and the sum runs over the five light (massless) quarks in the loop.

The matrix element for the tree-level process $\chi\chi \rightarrow q\bar{q}g$ is

$$\begin{aligned} \mathcal{M}_{\text{tree}} = & \frac{g_s}{\sqrt{2}} i\varepsilon^{\mu\nu\alpha\beta} \varepsilon_\mu^{*c} q_{3\alpha} (q_1 + q_2)_\beta \left\{ \frac{1}{2} \left[\frac{|g_r|^2}{M_{\tilde{q}_R}^4} + \frac{|g_\ell|^2}{M_{\tilde{q}_L}^4} \right] \bar{u}(q_1) T^c \gamma_\nu v(q_2) \right. \\ & \left. + \frac{1}{2} \left[\frac{|g_r|^2}{M_{\tilde{q}_R}^4} - \frac{|g_\ell|^2}{M_{\tilde{q}_L}^4} \right] \bar{u}(q_1) T^c \gamma_\nu \gamma_5 v(q_2) \right\}, \end{aligned} \quad (9)$$

where we neglect terms subleading in m_χ^2/M_q^2 . Only the vectorlike part of $\mathcal{M}_{\text{tree}}$ will interfere with the dimension-six amplitude. Summing over final-state polarizations and colors and averaging over initial-state polarizations, we obtain the interference term [2],

$$\frac{1}{4} \sum_{\text{pols}} 2 \text{Re} [\mathcal{M}_{\text{tree}} \mathcal{M}_{\text{split}}^*] = -4\alpha_s^2 \sum_{q'} \left[\frac{|g_r|^2}{M_{\tilde{q}_R}^2} + \frac{|g_\ell|^2}{M_{\tilde{q}_L}^2} \right] \sum_q \left[\frac{|g_r|^2}{M_{\tilde{q}_R}^4} + \frac{|g_\ell|^2}{M_{\tilde{q}_L}^4} \right] [(q_1 \cdot q_3)^2 + (q_2 \cdot q_3)^2], \quad (10)$$

where we have summed over the five light internal quarks q' in $\mathcal{M}_{\text{split}}$ of Eq. (8) and over the five light external quarks q . Integrating over the phase space, we find the contribution to the cross section from the interference term [2],

$$v_{\text{rel}} \sigma_{\text{int}} = -\frac{\alpha_s^2 m_\chi^2}{32\pi^3} \frac{2m_\chi^2}{3} \sum_{q'} \left[\frac{|g_r|^2}{M_{\tilde{q}_R}^2} + \frac{|g_\ell|^2}{M_{\tilde{q}_L}^2} \right] \sum_q \left[\frac{|g_r|^2}{M_{\tilde{q}_R}^4} + \frac{|g_\ell|^2}{M_{\tilde{q}_L}^4} \right]. \quad (11)$$

The form of this interference term is rather similar to that of the dimension-six cross section given in Eq. (5). In the special case of degenerate right- and left-handed squarks, $M_{\tilde{q}_R} = M_{\tilde{q}_L} \equiv M_{\tilde{q}}$, we find that the interference term is related to the dimension-six cross section by a multiplicative constant: $v_{\text{rel}}\sigma_{\text{int}} = [-2m_\chi^2/3M_{\tilde{q}}^2]v_{\text{rel}}\sigma_{\text{dim6}}$. Combining this result with the NLO corrections to $\chi\chi \rightarrow gg$ with $N_f = 5$ light flavors and a renormalization scale $\mu = 2m_\chi$, we obtain

$$v_{\text{rel}}\sigma = v_{\text{rel}}\sigma_{\text{dim6}} \left[1 + \frac{221}{12} \frac{\alpha_s^{(5)}(2m_\chi)}{\pi} - \frac{2}{3} \frac{m_\chi^2}{M_{\tilde{q}}^2} \right]. \quad (12)$$

For $m_\chi \sim 100$ GeV the NLO correction is roughly 60%; for, e.g., $M_{\tilde{q}} \sim 2m_\chi$, the interference term cancels off about one quarter of the NLO correction. We note that the expansion of the squark propagator in $\chi\chi \rightarrow gg$ yields correction terms starting at order $m_\chi^4/M_{\tilde{q}}^4$ in the square brackets in Eq. (12); these are beyond our present interest.

Outlook: In the region of SUSY parameter space where virtual squark exchange gives a sizable contribution to neutralino annihilation, the neutralinos are “p-wave annihilators” during freeze-out in the early universe; i.e., they annihilate through the velocity-dependent $\chi\chi \rightarrow q\bar{q}$ process, which is not suppressed by the quark mass. In this situation, the process $\chi\chi \rightarrow gg$ is subdominant and our QCD corrections constitute only about 1 – 2% of the dominant cross section. They are thus of little importance for computing the relic neutralino abundance. However, for neutralino annihilation at the present time the relative neutralino velocity is much lower, leading to a much smaller tree-level $\chi\chi \rightarrow q\bar{q}$ cross section. In this situation, the $\chi\chi \rightarrow gg$ cross section can be as large or larger than $\chi\chi \rightarrow q\bar{q}$, so that the QCD corrections computed here could have a significant impact on the computation of gamma ray and neutrino fluxes from neutralino annihilation in the galactic halo and inside the sun, respectively. In [2] we provided the spin-averaged squares of matrix elements for the dimension-six, dimension-eight, and interference term contributions to $\chi\chi \rightarrow q\bar{q}g$ in terms of the final-state particle momenta to facilitate their implementation into Monte-Carlo generators for the computation of indirect dark matter detection rates.

I thank Vernon Barger, Wai-Yee Keung, Gabe Shaughnessy, and Adam Tregre for an enjoyable and fruitful collaboration that led to these results. This work was supported by the Natural Sciences and Engineering Research Council of Canada.

REFERENCES

1. V. Barger, W. Y. Keung, H. E. Logan, G. Shaughnessy and A. Tregre, Phys. Lett. B **633**, 98 (2006).
2. V. Barger, W. Y. Keung, H. E. Logan and G. Shaughnessy, Phys. Rev. D **74**, 075005 (2006).
3. P. Gondolo, J. Edsjo, P. Ullio, L. Bergstrom, M. Schelke and E. A. Baltz, JCAP **0407**, 008 (2004).
4. S. L. Adler, “Perturbation Theory Anomalies,” in *Lectures on Elementary Particles and Quantum Field Theory*, Vol. 1, ed. S. Deser, M. Grisaru and H. Pendleton (M.I.T. Press, Cambridge, MA), pp. 3–164 (1970).
5. S. L. Adler and W. A. Bardeen, Phys. Rev. **182**, 1517 (1969).
6. M. Spira, A. Djouadi, D. Graudenz and P. M. Zerwas, Nucl. Phys. B **453**, 17 (1995).